

Nutrition and Food Technology: Open Access

Research Article

Volume: 2.1

Open Access

Prevalence of Coliforms, Antibiotic Resistant Coliforms and *E. Coli* Serotypes in Raw Milk and Some Varieties of Raw Milk Cheese in Egypt

Abd El Rahaman M Elbagory^{1*}, Ahmed M Hammad¹, Alzahraa² and MA Shiha²¹Department of Food Hygiene and Control, Faculty of Veterinary Medicine University of Sadat City, Egypt
² Menofia University Hospital, Shibin Elkom, Egypt***Corresponding author:** Abd El Rahaman M Elbagory, Department of Food Hygiene and Control, Faculty of Veterinary Medicine University of Sadat City, Egypt, **E-mail:** Elbagory200@yahoo.com**Received date:** 13 Jan 2016; **Accepted date:** 02 Feb 2016; **Published date:** 08 Feb 2016.**Citation:** Elbagory AM, Hammad AM, Alzahraa, Shiha MA (2016) Prevalence of Coliforms, Antibiotic Resistant Coliforms and *E. Coli* Serotypes in Raw Milk and Some Varieties of Raw Milk Cheese in Egypt. *Nutr Food Technol* 2(1): doi <http://dx.doi.org/10.16966/2470-6086.114>**Copyright:** © 2016 Elbagory AM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The present study investigated the prevalence of coliforms, antibiotic resistant coliforms and *E. coli* serotypes in raw milk and some varieties of raw milk cheese in Egypt. One Hundred and twenty samples of raw milk and white soft cheese (salted and unsalted) and hard cheese (Ras) collected from different areas in El Minofia governorate, Egypt were examined. The obtained results declared that coliform members including *E. coli*, *Citrobacter diversus*, *C. freundii*, *Enterobacter aerogenes*, *E. agglomerans*, *E. cloacae*, *Klebsiella pneumoniae* sub. spp. *ozaenae* and *K. pneumoniae* sub. spp. *pneumoniae* were isolated at varying percentage from examined samples. Kirby-Bauer disk diffusion test was used to examine bacterial susceptibility to ten antimicrobial agents including chloramphenicol, impenem, sulbactam+cefoperazone, ampicillin+sulbactam, amoxicillin+clavulanic acid, cefotaxime, gentamycin, amikacin, cefoperazone and tetracycline. Some strains showed multidrug resistant phenotype. The previously tested coliform strains, in order, showed resistance to AMC ; SAM & AMC ; AMC & CFP ; AMC & TE and C, AMC, CFP & TE antibiotics, while each showed sensitivity to the other tested antibiotics. The isolated *E. coli* proved to belong to 5 serotypes; O₁₁₁:K₅₈ (B₉) EHEC, O₁₂₄:K₇₂ (B₁₇) EIEC, O₁₁₉:K₅₉ (B₁₀) EPEC, O₅₅:K₅₉ (B₅) EPEC and untypable strains. In conclusion, this study provides meaningful data on the incidence of coliforms in raw milk cheese in Egypt and merely underlines the need for implementation of strict hygiene measures in preparation of raw milk cheese in Egypt. The most alarming trend detected in this study was the determination of antibiotic resistant and potentially toxigenic *E. coli* strains. This discovery makes ongoing surveillance of coliforms a high priority to identify emerging harmful strains within food production.

Keywords: Coliforms; Antibiotic resistant coliforms; *E. coli* serotypes; Cheese

Introduction

Although milk has an outstanding nutritional quality, yet it is also an efficient vehicle for transmission of food borne pathogens that pose a serious threat to human health, and constitute about 90% of all dairy related diseases [1]. Coliforms particularly *E. coli* are frequently used in the microbiological analysis of food as an indicator of poor hygienic condition. Moreover, coliform test is used to measure the quality of practices used to minimize microbial contamination of dairy products and as an approved safety indicator in HACCP system [2,3]. Contamination of cheese with coliforms especially fecal coliforms gives an indication of either direct or indirect fecal contamination and considered as a mirror for the degree of disregard of numerous hygienic rules during processing and marketing [4]. Some members of coliforms are responsible for the development of objectionable taints in milk and its products rendering them of inferior quality or even unmarketable, moreover certain serotypes of *E. coli* are associated with gastroenteritis and food poisoning outbreaks [2].

Antibiotic resistant coliforms have increased worldwide that leading to failures in treatment of human infectious diseases and they are a major concern in the anti-infective therapy of both humans and animals. Moreover they are considered dangerous healthy, social and economic problem in the world where antibiotic resistance of coliforms have a biological danger, which increases the disease in animals and human [5]. Therefore the present study is planned to isolate and identify coliforms, serotyping of isolated *E. coli* and determination of antibiotic resistance for various members of isolated coliforms.

Materials and Methods

Collection and preparation of cheese samples

One hundred and twenty samples; 30 each of raw milk, white salted soft, white unsalted soft cheese and Ras cheese were collected from dairy shops from different places in Menofia province, Egypt in clean sterile polyethylene bags and transferred in ice box to the laboratory with a minimum of delay to be examined. Each cheese sample was thoroughly mashed, then 25 grams from each prepared sample was aseptically added into 225 ml of 2% Sodium citrate and thoroughly mixed till completely emulsified.

Isolation and identification of coliforms

The technique recorded by APHA [6] was used for isolation of coliforms from prepared raw milk and prepared cheese samples. Isolated purified colonies of coliforms were identified morphologically and biochemically according to Holt et al. [7].

Serological identification of *E. coli* isolates [4]

Isolated strains of *E. coli* were identified serologically, using polyvalent and corresponding monovalent antisera manufactured by Denka Seiken Ltd. Japan or Oxoid Ltd. where two separate drops of sterile distilled water were put on a clean glass slide and a portion of the colony from the suspected culture was emulsified with sterile loop in each drop of sterile distilled water on the slide to give a smooth fairly dense suspension. After that, to one of the two suspensions, one drop of sterile distilled water

was added and mixed to be used as negative control while one drop of polyvalent antiserum was added to the other one and titled backward and forward up to one minute. Agglutination was observed when a strongly positive agglutination with one of pools of polyvalent serum. A further portion of the colony was inoculated into nutrient agar slant and incubated for further growth to get a culture for testing with monovalent sera. A heavy suspension of organisms from each slope culture was prepared in distilled water and slide agglutination test was performed with O and K sera to identify the O and K antigens. *E. coli* polyvalent 2, 3 and 4 and *E. coli* monovalent antisera; O26: K60 (B6), O44: K74 (B-), O55: K59 (B5), O78: K80 (B) O86: K (B7), O11: K58 (B9), O112: K66 (B1), O114: K9, O119: K69 (B19), O124: K72 (B17), O125: K70 (B15), O126: K71 (B16), O127: K63 (B8) O128: K67 (B12), O142: K86 and O157:H7 were used.

Antibiogram of some isolated coliforms

Characterized strains were tested for their susceptibility to 10 antimicrobial agents by the disc diffusion method according to the guidelines of the Clinical and Laboratory Standards Institute [8]. The antimicrobial agents tested were chloramphenicol (30 µg), imipenem (10 µg), sulbactam+cefoperazone (105 µg), ampicillin+sulbactam (20 µg), amoxicillin+clavulanic acid (30 µg), cefotaxime (30 µg), gentamycin (10 µg), amikacin (30 µg), cefoperazone (75 µg) and tetracycline (30 µg). The zone diameter for individual antimicrobial agents was then translated into susceptible, intermediate and resistant categories according to CLSI [8].

Discussion

Isolated coliforms

Data in figure 1 showed that the incidence of the isolated coliforms in the examined raw milk samples were *E. coli* (7/30, 23.33%), *Citrobacter diversus* (7/30, 23.33%), *C. freundii* (8/30, 26.67%), *Enterobacter aerogenes* (7/30, 23.33%), *E. agglomerans* (3/30, 10%), *E. cloacae* (4/30, 13.35%), *Klebsiella pneumoniae* sub. spp. *ozaenae* (6/30, 20%) and *K. pneumoniae* sub. spp. *pneumoniae* (4/30, 13.33%). While the isolated coliforms in the examined salted white soft cheese samples were *E. coli* (4/30, 13.3%), *Citrobacter diversus* (4/30, 13.33%), *C. freundii* (6/30, 20%), *Enterobacter aerogenes* (4/30, 13.33%), *E. agglomerans* (2/30, 6.67%), *E. cloacae* (6/30, 20%), *Klebsiella pneumoniae* sub. spp. *ozaenae* (3/30, 10%) and *K. pneumoniae* sub. spp. *pneumoniae* (6/30, 20%). Coliforms other than *E. coli* could be isolated with different percentage by Digrak & Ozcelik and Amer et al. [9,10]. The incidence of *E. coli* substantiated what have been recorded by Nichols et al. and El-Gamal & Abdel-Khalek [11,12]. It has been demonstrated that *K. pneumoniae* is an important cause of nosocomial infections like pneumonia,

septicaemia, urinary tract and soft tissue infections [13,14]. *K. oxytoca* was reported as enterotoxigenic microorganism and cause an antibiotic associated hemorrhagic colitis [15]. Some species of genus *Enterobacter* as *E. zakazakii*, unlike other enteric organisms, may cause highly lethal syndrome of bacteremia and meningitis with central nervous involvement in neonates [16]. Moreover the horizontal transfer of Shiga-like toxin genes to some species of genus *Citrobacter* as *C. freundii* which was responsible for an outbreak among school children in Germany, further highlight the public health importance of coliforms [17]. Data in figure 1 showed that the incidence of the isolated coliforms in the examined unsalted soft cheese were *E. coli* (4/30, 13.3%), *C. diversus* (5/30, 16.67%), *C. freundii* (7/30, 23.33%), *E. aerogenes* (6/30, 20%), *E. agglomerans* (2/30, 6.67%), *E. cloacae*, (2/30, 6.67%), *K. pneumoniae* sub. spp. *ozaenae* (5/30, 16.67%) and *K. pneumoniae* sub. spp. *pneumoniae* (3/30, 10%). Coliforms other than *E. coli* could be recorded by Moustafa, Zeinhom and Ahmed [18,19,20]. The incidence of *E. coli* in examined samples agree to some extent to what have been recorded by Hamid & El Owni [21]. The high rate of contamination in this type of cheese may be due to handling, transportation or storage in a place contaminated with these types of bacteria or during processing of cheese. This type contain no salt, where salting process is important in preservation and as a flavor enhancer, reducing the moisture content in the cheese [22] and suppress the growth of undesirable microorganisms [23].

Data in figure 1 showed that the incidence of the isolated coliforms in the examined hard cheese (Ras or Roomy cheese) were *E. coli* (3/30, 10%), *Citrobacter diversus* (2/30, 6.67%), *C. freundii* (3/30, 10%), *Enterobacter aerogenes* (4/30, 13.34%), *E. agglomerans* (1/30, 3.33%), *E. cloacae* (3/30, 10%), *Klebsiella pneumoniae* sub. spp. *ozaenae* (4/30, 13.33%) and *K. pneumoniae* sub. spp. *pneumoniae* (2/30, 6.67%). Similar incidence of *E. coli* in examined samples was recorded by Elessawy [24]. Similar coliforms other than *E. coli* could be recorded by Mohamed, Hassan and Sadek et al. [25-27]. The stages of processing of Ras cheese may minimize the risk of contamination and result in low level of contamination, but may be post processing as during transportation, storage or handling by unclean handlers [28].

Antibiotic resistant coliforms:

The results obtained in figures (2-6) explained the antibiotic resistance to coliform isolates including *E. coli* that isolated from milk and cheese samples. The tested coliform strains, in order *C. diversus*, *C. freundii*, *E. cloacae*, *K. pneumoniae* sub. spp. *pneumoniae* and *E. coli* O₁₁₁ showed resistance to AMC; SAM & AMC; AMC & CFP; AMC&TE and C, AMC, CFP&TE antibiotics, while each showed sensitivity and intermediate resistance to the other tested antibiotics. Wang et al. [29] pointed out that the most examined *C. freundii* isolates were resistant to gentamicin,




Figure 1: Incidence of coliform organisms in different cheese samples




Figure 2: Susceptibility of *C. diversus* to certain types of antibiotics





Figure 3: Susceptibility of *C. freundii* to certain types of antibiotic

Figure 6: Susceptibility of *E. coli* O₁₁₁:K₅₈ to certain types of antibiotics




Figure 4: Susceptibility of *E. cloacae* to certain types of antibiotic




Figure 5: Susceptibility of *K. pneumoniae* sub. spp. *pneumoniae* to certain types of antibiotics

Klebsiella spp. isolates showed susceptibility to imipenem followed by ciprofloxacin, piperacillin/tazobactam combination and ciftazidime. The imipenem antibiotic to which *Klebsiella* spp. was highly sensitive and is a drug of choice. Nam et al. [33] recorded that *C. freundii*, *E. cloacae* and *Klebsiella pneumoniae* isolates have strong antimicrobial activity against amikacin, gentamicin, and piperacillin whereas rifampin, cephalothin, ceftazidime, and ampicillin were ineffective against most of the tested bacterial species. Paneto et al. [34] surveyed the occurrence of toxigenic *E. coli* isolated from raw milk cheese samples and tested its susceptibility to certain antimicrobial agents and found that the most frequent resistance was observed to the cephalothin, nalidixic acid, doxycycline, tetracycline and ampicillin, but all tested *E. coli* isolates from raw milk samples exhibited 100% resistance to rifampin and tetracycline and 50% resistance to nalidixic acid but were 100% sensitive against imipenem as recorded by Aftab Uddin et al. [35]. Sallam [36] recorded that *E. coli* isolates were tested for susceptibility to eleven antibiotics; ampicillin, amoxicillin, amoxicillin/clavulanic acid, oxytetracycline, tetracycline, doxycycline, ciprofloxacin, enrofloxacin, neomycin, gentamicin and erythromycin. The resistance to such antibiotics were 50%, 50%, 7%, 67%, 67%, 40%, 23%, 23%, 23%, 13%, and 100%, respectively. It is evident from the last few years that some strains of *Klebsiella*, *Enterobacter* and *Citrobacter* spp. had been isolated from stools and the intestinal contents of both children and adults in several epidemiological studies of acute and chronic disturbances. Moreover, certain members of *Citrobacter* spp. had been suspected to cause enteric infection. *C. freundii* had been found among urinary and other pyogenic infections in humans [37]. Also, there is a loose association between *E. coli* and possible enteric pathogens [10]. *Klebsiella pneumoniae* occurs as a commensal in the upper respiratory tract and in the intestine and may be associated with catarrhal conditions in the respiratory tract, paranasal sinusitis, conjunctivitis and pneumonia in man [38].

Serological typing of isolated *E. coli* strains:

It's evident from the results recorded in table 1 that some isolated *E. coli* strains could be serologically typed into serotypes O₁₁₁ (3 strains), O₁₁₉ (2 strains), O₅₅ (one strain) and O₁₂₄ (one strain). These findings substantiated what have been found by Saudi and Moawad, Ahmed and Sallam and El-Shinawy et al. [39-41]. *E. coli* has been established to be among the etiological agents causing enteritis and several extra gastrointestinal diseases. Also, the organisms are recognized as pathogens for human and ruminants [42]. In recent years, much attention has been paid towards *E. coli* because of its importance as an organism of true fecal origin with the possible existence of associated enteric pathogens. Also, the public health hazard of enteropathogenic *E. coli* has been emphasized by many investigators

tobramycin, and aztreonam. The results indicated that the susceptible rates of *C. freundii* to aminoglycosides and ciprofloxacin are decreased markedly. Yagoub et al. [30] declared that the examined *Enterobacter* spp. isolated from raw milk samples had a wide range of resistance to penicillin, clindamycin, amoxicillin and ampicillin, while Farzana et al. [31] showed that the tested *Enterobacter* isolate was sensitive to nalidixic acid, tetracycline and showed little resistance to chloramphenicol. *Klebsiella* spp. isolate was resistant to nalidixic acid and little to chloramphenicol as have been showed by Farzana et al. [31], while Chauhan et al. [32] illustrated that the tested

Examined samples	No of sample	Isolated <i>E. coli</i>	Strain Character*
Raw milk	1	O ₁₁₁ :K ₅₈ (B ₉)	EHEC
	1	O ₁₂₄ :K ₇₂ (B ₁₇)	EC
	1	O ₁₁₉ :K ₆₉ (B ₁₉)	EPEC
White salted cheese	1	O ₁₁₁ :K ₅₈ (B ₉)	EHEC
	1	O ₁₁₉ :K ₆₉ (B ₁₉)	EPEC
White unsalted cheese	1	O ₅₅ :k ₅₉ (B ₅)	
	1	Untypable	-
Ras cheese	1	O ₁₁₁ :K ₅₈ (B ₉)	EHEC

Table 1: Serological identification of isolated *E. coli* of examined samples.

as they have been implicated in human cases of gastroenteritis, epidemic diarrhea in infants, sporadic diarrhea in children as well as in cases of food poisoning [43]. However 6 main types of diarrheagenic *E. coli* have been associated with food borne disease; enterotoxigenic *E. coli* (ETEC) which is responsible for watery diarrhea (travelers' diarrhea) as consequence to production of heat labile toxin (Cholera like toxin) and heat stable toxin (diarrheal toxin). Enteropathogenic *E. coli* (EPEC) which is associated with infantile diarrhea, Enterohaemorrhagic *E. coli* which may result in bloody diarrhoea, hemorrhagic colitis, hemolytic uremic syndrome and thrombocytopenic purpura. Enteroaggregative *E. coli* (EaggEC) which aligns themselves in parallel rows on either tissue cells or glass described as 'stacked brick-like' then elaborate heat labile toxin, antigenically related to haemolysin and a plasmid encoded that stables toxin resulting in persistent watery diarrhea especially in children. Enteroinvasive *E. coli* (EIEC) which may result in fever and profuse watery diarrhoea containing mucous and streaks of blood. Diffusely adherent *E. coli* (DAEC), have been associated with diarrhea in some studies but not consistently [44].

In conclusion, this study provides meaningful data on the incidence of coliforms in raw milk cheese in Egypt and merely underlines the need for implementation of strict hygiene measures in preparation of raw milk cheese in Egypt. The most alarming trend detected in this study was the determination of antibiotic resistant and potentially toxigenic *E. coli* strains. This discovery makes ongoing surveillance of coliforms a high priority to identify emerging harmful strains within food production.

References

- Ryser ET (1998) Public Health Concerns. In: Marth EH, Steele J (eds) Applied Dairy Microbiology. 2nd Edition, Merrell Dekker Inc., New York 263-404.
- Banwart GJ (1989) Basic Food Microbiology. 2nd edition, Vien Nostrand Reinhold Publisher, New York.
- Jay JM (2000) Modern food microbiology. 6th edition, Van Nostrand Reinhold Company, New York.
- Varnam AH, Evans MG (1991) Food borne pathogens: An illustrated text. Chapter 13, Book House, Tomington palace, England. 267.
- European Food Safety Authority (2008) Foodborne antimicrobial resistance as a biological hazard. Scientific Opinion of the Panel on Biological Hazards (Question No EFSA–Q-2007- 089).
- APHA (2004) Standard Methods for the Examination of Dairy Products. In: Wehr HM, Frank JH (eds) 17th edition, Washington, DC, USA.
- Holt JG, Krieg NR, Sneath PHA, Staley JT, Williams ST (1994) Bergey's Manual™ of determinative bacteriology. 9th Edition, Williams &Wilkins, Baltimore.
- Clinical and Laboratory Standards Institute (2005) Performance standards for antimicrobial susceptibility testing. 15th informational supplement. CLSI document M100-S15, Clinical and Laboratory Standards Institute, Wayne, PA.
- Digrak M, Ozcelik S (1996) The identification of the coliform group bacteria isolated from the cheese marketed in Elazig. GIDA 21: 3-7.
- Amer IH, Abd El-Aal SFA, Kamal RM (2005) Comparative microbiological study on soft cheese manufactured by small or large scale production in Sharkia province, Egypt. Zagazig Vet J 33: 190-196.
- Nicholas G, Greenwood M, Louvois J, De-Louvois J (1996) The microbiological quality of soft cheese. PHLS-Microbiology-Digest 13: 68-75.
- El-Gamal AM, Abdel-Khalek A (1997) Quality control of white soft cheese in El-Dakahlia province, Egypt. Alex J Vet Sci 13: 784-790.
- Maslow JN, Brecher SM, Adams KS, Durbin A, Loring S, et al. (1993) Relationship between indole production and differentiation of klebsiella species: Indole positive and negative isolates of *Klebsiella* determined to be clonal. J Clin Microbiol 31: 2000-2003.
- Podschun R, Ulmann U (1998) *Klebsiella* spp. as nosocomial pathogens: Epidemiology, taxonomy, typing method and pathogenicity factors. Clin Microbiol Rev 11: 589-603.
- Minami J, Katayama S, Matsushita O, Sakamoto H, Okabe A (1994) Enterotoxin activity of *Klebsiella oxytoca* cytotoxin in rabbit intestinal loops. Infect Immun 62: 172-177.
- Gallagher PG (1990) Enterobacter bacteremia in pediatric patients. Rev Infect Dis 12: 808-1812.
- Neil MA (1997) Overview of veotoxigenic *E. coli*. J Food Prot 60: 1444-1446.
- Moustafa MM (2011) Hygienic status of milk and some dairy products received at university hostels in kaliobia governorate. Ph.D Thesis, Benha University, Egypt.
- Zeinhom MMA (2011) Monitoring of enteric pathogens in milk and some dairy products with special reference for Enterobacter sakazaki and *E. coli* O₁₅₇:H₇. Ph.D. Thesis, Department of Food Hygiene, University of Beni-Suef, Egypt.
- Ahmed EGH (2012) Microbiological evaluation of soft chesses sold in Beni-Suef markets. Thesis Ph.D. Thesis, Department of Food Hygiene, University of Beni-Suef, Egypt.
- Hamid OIA, El Owni OAO (2007) Microbiological properties and sensory characteristics of White Cheese (Gibna bayda) collected in Zalingei Area, West Darfur State. Res J Anim Vet Sci 2: 61-65.
- He FJ, Macgregor GA (2007) Dietary salt, high blood pressure and other harmful effects on health. In: Kilcast D, Angus F (eds) Reducing salt in foods: Practical strategies. Cambridge, UK 18-54.
- Fennema OR (1996) Food chemistry. 3rd Edition, Marcel Dekker, New York.
- Elessawy HNE (2000) Prevalence of Enteropathogenic *E. coli* in raw milk and some dairy products with special reference to *E. coli* O₁₅₇:H₇. M.Sc. Thesis, Cairo University.
- Mohamed JII (2004) Biogenic amines and their forming organisms in cheese. Ph. D. Thesis, Suez canal University, Egypt.
- Hassan MAA (2008) Assessment of probable microbiological hazards in some dairy products. M.Sc. Thesis Ain-Shams University.

27. Sadek ZI, Hosney IM, El-Kholy WI, El-Dairouty RK (2009) Comparative investigations for detection of foodborne microorganisms in Egyptian Hard Cheese (Ras) using conventional and fast biochemical tests. National Research Centre- Dokki; Giza, Egypt. *Global Veterinaria* 3: 189-195.
28. Zeinab I, Hosny IM, El-Kholy WI, El-Dairouty RK (2009) Comparative investigations for detection of foodborne microorganisms in Egyptian hard cheese 'ras' using conventional and fast biochemical tests. *Global Veterinaria* 3: 189-195.
29. Wang JT, Chang SC, Chen YC, Luh KT (2000) Comparison of antimicrobial susceptibility of *Citrobacter freundii* isolates in two different time periods. *J Microbiol Immunol Infect* 33: 258-262.
30. Yagoub SO, Awadalla NE, El-Zobeir ILM (2005) Incidence of some potential pathogens in raw milk in north Khartoum, Sudan and their susceptibility to antimicrobial agents. *J Anim Vet Adv* 4: 356-359.
31. Farzana K, Jabeen F, Shah SN (2005) Antibiotic resistance pattern among gram-negative bacteria isolated from raw milk in Multan. *Pak J Pharm Sci* 18: 11-16.
32. Chauhan S, Farooq U, Singh V, Kumar A (2013) Determination of prevalence and antimicrobial activity of ESBL (extended spectrum beta-lactamases) producing *klebsiella* species isolated from raw milk of Doon valley in India. *Int J Pharm Bio Sci* 4: 417-423.
33. Nam H, Lim S, Kim J, Joo Y, Jang K, et al. (2010) *In vitro* activities of antimicrobials against six important species of gram-negative bacteria isolated from raw milk samples in Korea. *Foodborne Pathog Dis* 7: 221-224.
34. Paneto BR, Schocken-Iturrino RP, Macedo C, Santo E, Marin JM (2007) Occurrence of toxigenic *E. coli* in raw milk cheese in Brazil. *Arq Bras Med Vet Zootec* 59: 2.
35. Uddin MdA, Motazzim-ul-Haque HMd, Noor R (2011) Isolation and identification of pathogenic *E. coli*, *Klebsiella spp.* and *Staphylococcus spp.* in raw milk samples collected from different areas of Dhaka city, Bangladesh. *Stamford J Microbiol* 1: 19-23.
36. Sallam AMA (2013) Studies on antimicrobial resistance and its relation to antibiotic residues. M.Sc. Thesis Cairo University, Egypt.
37. Abd El-Fatah EN (2007) Sanitary studies on fermented milks marketed at Zagazig markets. M.V.Sc. Thesis, FacVet Med Zagazig University, Egypt.
38. Abdel-Aziz A (1973) Medical Microbiology. Cairo University, Sherif Press, Cairo.
39. Saudi AMM, Moawad AA (1990) Incidence of Enterobacteriaceae in market milk in Cairo and its suburbs. *Assiut Vet Med J* 24: 134-139.
40. Ahmed AM, Sallam SS (1991) Public health significance of coliforms in raw milk and Domiaty cheese. *Assiut Vet Med J* 25: 50.
41. El-Shinawy SH, El-Kholy AM, El-Gwad MH (1995) Sanitary condition of market raw milk in Beni-Suef governorate, Beni-Suef. *Vet Med J* 5: 330-338.
42. Robertson DC (1988) Pathogenesis and Enterotoxins of diarrhoeagenic *E. coli*. In: Roth JA (ed) Virulence mechanisms of bacterial pathogens. American society for Microbiology, Washington, D.C.
43. Mahran MA (2000) Sanitary condition of goat's milk in Upper Egypt. M.V.Sc. Thesis, Faculty of Veterinary Medicine, Assiut University, Egypt.
44. Forsythe SJ (2000) The microbiology of safe food. 1st Edition, Blackwell Science Ltd, London 285.