

Mediators Involved in Obesity and Modulation of Food Intake by Antioxidants

Bruna Vidal Dias¹, Helena Dória Ribeiro de Andrade Previato² and Ana Carolina Pinheiro Volp^{1,3*}

¹Federal University of Ouro Preto (UFOP), Ouro Preto- Minas Gerais, Brazil

²State University of Campinas (UNICAMP), Campinas-São Paulo, Brazil

³Federal University of Mato Grosso (UFMT), Cuiabá-Mato Grosso, Brazil

*Corresponding author: Ana Carolina Pinheiro Volp, Federal University of Mato Grosso (UFMT), Cuiabá-Mato Grosso, Brazil, E-mail: anavolp@gmail.com

Received: 15 Nov, 2018 | Accepted: 22 Dec, 2018 | Published: 30 Dec, 2018

Citation: Dias BV, Previato HDRA, Volp ACP (2018) Mediators Involved in Obesity and Modulation of Food Intake by Antioxidants. *Nutr Food Technol Open Access* 4(1): dx.doi.org/10.16966/2470-6086.153

Copyright: © 2018 Dias BV, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: Obesity is a disease that involves adiposity excess associated to diverse factors, including behavior ones, such as hormonal, adipocytarious, neural and intestinal. The excess of adipose tissue results in inflammation, which begins with secretion of pro-inflammatory adipokines, activating inflammatory T cells and resulting in imbalance between antioxidants and oxidants compounds. The recuperation of this defense system may occur through endogenous or dietetics origin. Diet rich in antioxidants is inversely bound to metabolic disturbances. Therefore, these compounds may be regarded coadjutant potentials in the prevention of obesity.

Objectives: To review and to summarize critically studies related to themes concerning antioxidants and food intake mediators.

Methods: It was done a narrative review about food intake mediators in humans related to obesity and antioxidants intake.

Results: Overweight individuals possess an imbalance between intrinsic signals related to appetite and a nutritionally balanced diet is fundamental to oppose obesity. The intake of food rich in antioxidant may be a strategy to reach metabolic homeostasis and the reduction of damage caused by obesity.

Conclusion: The antioxidant compounds present an anti-obesity potential effect due to food intake mediators' modulation for opposing damage caused by excess of adipose tissue deposition.

Keywords: Food intake mediators; Obesity; Antioxidants; Polyphenols; Anthocyanins

Introduction

Obesity is a complex disease, which involves the excessive adiposity accumulation associated to diverse factors both behavioral as hormonal, adipocytarious, neural and intestinal [1].

The excess of adipose tissue deposition secretes many bioactive substances involved in the regulation of insulin, arterial pressure and inflammatory estate such as leptin, adiponectin, InterLeukin (IL), Tumor Necrosis Factor alpha (TNF- α), among others [2]. A subclinical inflammation is observed subsequently to the adiposity excess which begin from the pro-inflammatory adipokine, such as leptin, which activates inflammatory T cells, causing a progressive inflammation [3].

On the subclinical chronic inflammation occurs an imbalance between antioxidant and oxidant compounds, which favors the installation of an oxidative stress process. The maintenance and recuperation of this defense system may occur through enzymatic or non-enzymatic systems, and this may be of an endogenous origin

(through dismutase superoxide, catalase, glutathione peroxidase) or dietetic through the intake of food rich in Vitamin A (β -caroteno), Vitamin C (ascorbic acid), Vitamin E (α -tocoferol), copper, zinc, magnesium, selenium, caratenoids (lycopene) and phytochemical (resveratrol, catechins, quercetins, phenolic acid and others). These substances may act both as directly, neutralizing the action of free radicals, as indirectly, activating enzymatic systems capable of neutralize them [4]. Thus, oxidative stress is an important link between the intake of a hypercaloric diet and chronic diseases, such as obesity [5].

Food intake is modulated mainly through biological mechanisms such as homeostatic, which refer to hormonal regulators that initiate hunger and satiation (such as neuropeptide Y and tyrosine tyrosine peptide) or sign adiposity levels (such as leptin, ghrelin and insulin), which act in the hypothalamic encephalic centers and in the brainstem, sending and receiving signals according to available energetic substratum in order to keep a proper energetic balance; or may not be homeostatic (hedonic), which refer to awarding cerebral

system. So palatable food (rich mainly in fat and sugar) are consumed even after the energetic requirement being accomplished. On the other hand, non-palatable food (no sugar and fat) are not desired on the same way by the brain [6,7].

The influence of biological factors and their alterations induced by diets on corporal composition are not totally clarified yet [8]. Gastrointestinal system presents special chemoreceptors and mechanoreceptors that monitor quantity and quality of food intake and nutrients. Therefore, the information arrives at the hypothalamus through afferent nerve fibers that understand the signals related to current necessities [7]. Many hormones are involved in these feedback mechanisms between gastrointestinal tract and brain, and those hormones may be liberated before or after food intake, whether they are orexigenic or anorexigenic, always looking for the balance between intakes, spent and stocked energy. Among the involved mediators in such regulation, it may be mentioned Peptide Tyrosine Tyrosine (PYY), Pancreatic Polypeptide (PP), alpha-Melanocyte Stimulating Hormone (alpha-MSH), Peptide C, chemerin and beta-endorphin [9].

The homeostasis of these mediators depends on individual factors such as corporal composition and lifestyle, including exercise and nourishment [10]. Thus, a diet rich in fruits and vegetables is inversely linked to metabolic disturbances [11]. Grains, teas and wines are rich in polyphenols, which are natural substances composed by phenolic rings. Polyphenols might be classified as stilbenes, lignans, phenolic acids and flavonoids [12].

Polyphenols present many benefits for health due to their antioxidant potential effect, anti-inflammatory [13] and insulin potentiator, causing a plasmatic glucose reduction in diabetic [14] and obese people [15]. However, their effects on the regulation of neuropeptides and neural-hormones involved at food intake are not totally clarified [16].

The objective of this paper was to review studies for a higher understanding about the effect of antioxidants on food intake mediators.

Methodology

This paper consists on a narrative review of studies about food intake mediators' and their relation with antioxidants consumption.

It was done a bibliographical search on databases *PubMed*, *SciElo* and *Science Direct*, considering the following keywords: "food intake mediators", in association to "obesity", "antioxidants", "polyphenols" and "anthocyanins" and their respective correspondents in English: "food intake mediators", "obesity", "antioxidants", "polyphenols", and "anthocyanins".

In order to assess the action of food intake mediators, it was included interventions done in humans, which would assess their action over metabolic parameters associated to obesity. It was included studies with humans and animals to assess the action of antioxidants on food intake parameters, considering that the literature on the topic is still scarce.

Results

Many proteins involved in the arterial pressure, fat and glucose metabolism regulation, such as leptin and chemerin are adipokine secreted by adipose tissue [17]. It is also known that many cytokines and proteins of acute stage related to subclinical inflammation are highly found in obese patients. An imbalance between pro and anti-inflammatory cytokines induces the inflammation estate, which occurs when there is an adipose tissue, forming a cycle responsible for

oxidative stress, until its cause is reverted [18].

The excessive intake of macronutrients is one of the biggest contributors for the increasing of obesity prevalence [19]. However, the act of eat excessively is not always something conscious or compulsive; studies show that there may be a defective answer of intrinsic signs related to appetite in overweight individuals [20-22].

The act of eating is highly related to many processes, which result in the energetic balance, such as hunger, satiation and repletion. Hunger is considered as the urge that leads to the action of eating; satiation is the period in which eating is interrupted and repletion is the period between meals, where there is an eating inhibition. This "food desire" may be, or not, due to a physiologic necessity, so it can be stimulated by hormonal signal due to lack of energetic substratum, such as olfactory or visual external stimuli [23].

The ingestion/expenditure of nutrients are controlled by neural structures, which involve specific neurochemical and neuroendocrine systems and, when it happens a commitment in these systems, occurs a consequent imbalance of eating behavior [24]. With this, there is a complex interaction between pre-absorbing, absorbing and post-absorbing stages, which determine food intake and energetic expenditure. Imbalanced intakes of macronutrients and micronutrients may induce an alteration of some mediators' production, which intervene directly on corporal composition, biochemical and clinical profile [25].

Thus, balanced diet is a fundamental strategy against obesity [26]. Potential mechanisms related to the benefits of a diet modification include: improvement of insulin resistance; improvement of insulin secretion depending on glucose at beta pancreatic cells; hunger/satiation system regulation through proper working of the mediators responsible for appetite and inflammation attenuation [27].

Many neuropeptides have been studied in order to better understand central factors related to ingestion control and obesity etiology [28,29,8].

Food intake mediators and obesity in humans

Many food intake mediators are responsible for energetic homeostasis and metabolic disturbances, which may proceed to an imbalance among food intake versus energetic expenditure [30].

PYY is an intestinal peptidic hormone secreted on post-prandial period proportionally to the ingested caloric quantity. This is secreted by L cells of distal portion of large and small intestine and it reduces intestinal mobility and, therefore, it improves satiation and reduces the intake of food [31].

Studies show that obese individuals present minor concentrations of endogen PYY in fasting and post-prandial period when compared to non-obese ones [32-34].

Brandão PP, et al. [35] assessed PYY concentration after and before meals in obese women with or without eating compulsion episodes. A total of 25 women from 32 to 50 years old were studied (9 eutrophic without eating compulsion episodes, 9 obese ones with eating compulsion episodes and 7 obese women without eating compulsion episodes). It was found difference on PYY levels when the thin women group were compared to the obese group (the obese ones presented lower levels). The group with eating compulsion episode presented the lowest levels of PYY, reinforcing its direct relation to the ingested caloric quantity and obesity.

In a study carried out to determine PYY effects managed peripherally over food intake and activation of enteric neurons and rhombencephalon found a reduction of 10% sucrose solution ingestion and activation of enteric neurons before the vagal dorsal complex, suggesting that these act peripherally before acting centrally [36].

Another determining factor of concentrations is the ingested caloric density, which directly influences the serum PYY concentration, thus presented an important relation to obesity genesis [37].

Another marker is the pancreatic polypeptide, a peptide hormone produced by special cells on pancreatic islet, called F cells, which, as with PYY, influence the energetic balance, for they reduce food intake [38]. This hormone may be produced also by exocrine way on distal intestine [39].

Studies showed that a higher secretion of PP is related to reduction of food intake and that circling levels of this hormone keep elevated up to 6 hours after the meal [40-42].

Low concentrations of PP were found concomitantly to the increase of NPY secretion (an orexigenic hormone) on the arched core hypothalamus on obese humans [43]. Edelsbrunner ME, et al. [44] observed that through knocking down the gene related to PP (gene Y4) occurs a higher food intake and a consequent gaining of weight in rats, confirming its anorexigenic action.

A study that assessed PP infusions effects done in the morning and afternoon in patients with Prader-Willi Syndrome (a genetic disease which leads to child obesity) observed a caloric ingestion reduction of approximately 12% [45]. With this, administration of PP in obese individuals might help to reduce the ingestion through ghrelin hormone concentration reduction (an orexigenic hormone) and thus control and restore the energetic balance [46].

Chemerin is a chemotactic adipokine, which connects to CMKLR1 receptor attached to the G protein that participates on the metabolic regulation of adipocytes, and, due to this its role on inflammation and metabolism may assist the understanding of this relation between obesity and chronic inflammation [47,48].

Besides many obese individuals present higher concentration of chemerin when compared to thin individuals, it was shown association of concentrations of this adipokine with triglycerides, arterial pressure, insulin resistance and corporal mass in non-diabetic individuals [48,49].

Schmid A, et al. [50] reported that oral ingestion of a lipidic emulsion, free from carbohydrates and proteins, regulates negatively the serum concentrations of chemerin in healthy humans. However, the molecular mechanisms of this regulation in humans are not clarified [50].

A good food intake mediator is the beta-endorphin, a peptide of melanocortins family found in organisms in lower concentrations under normal conditions, which may increase up to 10 times when in stress [51]. Studies show that central infusion of beta-endorphin stimulates food intake [52-54].

Considering that beta-endorphin is connected to the reward system under non-private conditions, it may be considered a good marker to assist the understanding of corporal composition and dietetic parameters, since its effects are related to palatable food intake and these are associated to overweight and obesity [55].

Alpha-MSH (hormone stimulating of alfa-melanocytes) also participates of the hedonic control of eating, as beta-endorphin, it is a neuropeptide of the melanocortins family [56]. It has an anorexigenic

action and participates of intrinsic processes related to metabolic process on obesity [57].

The anorexigenic action of alpha-MSH seems to occur due to the neurotrophic factor stimulation derived from the brain (BDNF), an endogen protein responsible for regular synapses stimulated both peripherally as centrally, after the activation of melanocortin receptor [58]. In humans, mutations on the gene of MC4R receptor may result in cases of obesity, which might appear in severe form even in the childhood [59].

The mRNA expression of its receptor possesses a direct relation to adipose tissue on obese individuals [60], and its related to the increase of lipolysis on white adipose tissue [61], muscular insulinic sensibility [62], thermogenesis on brown adipose tissue [63] and reduction of insulin excretion on pancreatic beta cells [64], thus characterizing a good correspondent of subclinical inflammation [60].

Food intake mediators and antioxidants

There are growing evidences on the protection role of health through fruits and vegetables consumption, which are the main contributors for polyphenols intake [65-67]. These compounds' ingestion has presented effects on the plasmatic glucose concentration reduction in obese individuals, with metabolic syndrome and type 2 diabetes [14,15]. These effects happen mainly for its antioxidant action. Studies showed that polyphenols distribute largely on the tissues where they act as exogenous antioxidants on neutralization of free radicals and consequent attenuation of oxidative stress [68,69].

While a small part of polyphenols may be absorbed on the small intestine, the bigger part ingested on the diet is not bioavailable on the distal portion of the human intestine, exerting its biologic activity on the intestinal tract through the interaction of colon microbiota [70].

These compounds may act directly on the encephalon. Studies in rats showed that they pass the hematoencephalic barrier after assessing the catechins concentration (a category of flavonoids) in encephalic tissue after polyphenol extract intake. However, necessary concentration for this was not totally clarified yet [71-73]. Despite that, the effects of neuropeptides and neuro-hormones regulation involved to food intake deserve attention, for it is a potential prevention and treatment of disturbances related to food intake such as obesity, diabetes, and others [16].

The intake of grape extract rich in polyphenol for 10 consecutive days by rats increased significantly the catechin concentrations and epicatechins (kinds of polyphenols) on encephalic tissue, result which was not observed on the group that received a single dose, which suggest that the long term intake may improve its action [74].

Studies proved that polyphenols intake causes a reduction of factors related to metabolic syndrome and obesity through the suppression of fat absorption on the intestine, thermogenesis activation, attenuation of chronic inflammation and fat oxidation [75-79,14].

Some studies in humans and in cultures of animal cells helped to clarify the potential role of polyphenols on neuro-hormones modulation related to food intake and energetic homeostasis [80].

The insulin, an important hormone that regulates the glucose levels on blood, has its action empowered with the polyphenols intake, causing an improvement on insulin resistance in obese patients with type 2 diabetes and/or metabolic syndrome [16].

Lu C, et al. [77] assessed the administration of a green tea in rats submitted to a diet rich in fat and observed a reduction on the ghrelin levels, a orexigenic hormone, caused by polyphenols contained in the

drink.

It was found by Weickert a potential anorexigenic effect caused by isoflavone intake (polyphenol contained in soy) due to the increasing of PYY concentration in healthy women [81].

Considering that inflammatory factors and oxidative estate are related to obesity, the reduction of those by antioxidant action through diet might favor the metabolic homeostasis [82-85].

Conclusion

Food intake mediators are very important to energetic homeostasis. Disturbances on the working of orexigenic or anorexigenic mediators may cause a metabolic dysfunction and, consequently, adipose tissue accumulation and obesity. Due to the capacity of reduce the inflammation damage and oxidative stress observed in obesity, the antioxidant compounds present potential anti-obesity effect due to the modulation of food intake mediators. However, more studies are necessary in humans to clarify this effect.

References

- Halpern ZSC, Rodrigues MDB, da Costa RF (2004) Physiological Determinants of Weight and Appetite Control. *Rev Psiquiatr Clin* 31: 150-153.
- Rosa G, Mello DB, Fortes MSR, Dantasa EHM (2013) Adipose tissue, metabolic hormones and exercise. *Andalusian Journal of Sports Medicine* 6: 78-84.
- Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, et al. (2009) CD8⁺ Effector T Cells Contribute to Macrophage Recruitment and Adipose Tissue Inflammation in Obesity. *Nat Med* 15: 914-920.
- Clarkson PM, Thompson HS (2000) Antioxidants: What Role Do They Play in Physical Activity and Health? *Am J Clin Nutr* 72: 637S-646S.
- Burneiko RCM, Diniz YS, Galhardi CM, Rodrigues HG, Ebaid GM, et al. (2006) Interaction of Hypercaloric Diet and Physical Exercise on Lipid Profile, Oxidative Stress and Antioxidant Defenses. *Food Chem Toxicol* 44: 1167-1172.
- Coll AP, Farooqi IS, O'Rahilly S (2007) The Hormonal Control of Food Intake. *Cell* 129: 251-262.
- Egecioglu E, Skibicka KP, Hansson C, Alvarez-Crespo M, Friberg PA, et al. (2011) Hedonic and Incentive Signals for Body Weight Control. *Rev Endocr Metab Disord* 12: 141-151.
- Ribeiro EB (2009) Studying the Central Control of Food Intake and Obesity in Rats. *Rev Nutr* 22: 163-171.
- Suzuki K, Simpson KA, Minnion JS, Shillito JC, Bloom SR (2010) The Role of Gut Hormones and the Hypothalamus in Appetite Regulation. *Endocr J* 57: 359-372.
- Wynne K, Stanley S, McGowan B, Bloom S (2005) Appetite Control. *J Endocrinol* 184: 291-318.
- Abete I, Goyenechea E, Zulet MA, Martínez JA (2011) Obesity and Metabolic Syndrome: Potential Benefit from Specific Nutritional Components. *Nutr Metab Cardiovasc Dis* 21: B1-B15.
- Manach C, Scalbert A, Morand C, Rémésy C, Jiménez L (2004) Polyphenols: Food Sources and Bioavailability. *Am J Clin Nutr* 79: 727-747.
- Landete JM (2012) Updated Knowledge about Polyphenols: Functions, Bioavailability, Metabolism, and Health. *Crit Rev Food Sci Nutr* 52: 936-948.
- Akilen R, Tsiami A, Devendra D, Robinson N (2012) Cinnamon in Glycaemic Control: Systematic Review and Meta Analysis. *Clin Nutr* 31: 609-615.
- Meydani M, Hasan ST (2010) Dietary Polyphenols and Obesity. *Nutr* 2: 737-751.
- Panickar K, Cao H, Qin B, Anderson RA (2009) Molecular Targets and Health Benefits of Cinnamon. In: Aggarwal BB, Kunnumakkara AB (eds) *Molecular Targets and Therapeutic Uses of Spices: Modern Uses for Ancient Medicine*. World Scientific 87-115.
- Trayhurn P (2007) Adipocyte Biology. *Obes Rev* 8: 41-44.
- Bullo M, Garcia-Lorda P, Megias I, Salas-Salvadó J (2003) Systemic Inflammation, Adipose Tissue Necrosis Factor, and Leptin Expression. *Obes Res* 11: 525-531.
- Carnell S, Kim Y, Pryor K (2012) Fat Brains, Greedy Genes and Parent Power: A Biobehavioural Risk Model of Child and Adult Obesity. *Inter Rev Psychiatry* 24: 189-199.
- Batterink L, Yokum S, Stice E (2010) Body Mass Correlates Inversely with Inhibitory Control in Response to Food among Adolescent Girls: an fMRI Study. *Neuroimage* 52: 1696-1703.
- Houben K, Nederkoorn C, Jansen A (2014) Eating on Impulse: The Relation Between Overweight and Food-Specific Inhibitory Control. *Obesity (Silver Spring)* 22: E6-E8.
- Nederkoorn C, Coelho JS, Guerrieri R, Houben K, Jansen A (2012) Specificity of the Failure to Inhibit Responses in Overweight Children. *Appetite* 59: 409-413.
- Blundell J, de Graaf C, Hulshof T, Jebb S, Livingstone B, et al. (2010) Appetite Control: Methodological Aspects of the Evaluation of Foods. *Obes Rev* 11: 251-270.
- Camilleri M (2006) Integrated Upper Gastrointestinal Response to Food Intake. *Gastroenterology* 131: 640-658.
- Grill HJ (2006) Distributed Neural Control of Energy Balance: Contributions from Hindbrain and Hypothalamus. *Obesity (Silver Spring)*: 216S-221S.
- Jacobs DR Jr, Gross MD, Tapsell LC (2009) Food Synergy: An Operational Concept for Understanding Nutrition. *Am J Clin Nutr* 89: 1543S-1548S.
- Ley SH, Hamdy O, Mohan V, Hu FB (2014) Prevention and Management of Type 2 Diabetes: Dietary Components and Nutritional Strategies. *Lancet* 383: 1999-2007.
- Pliquet RU, Fuhrer D, Falk S, Zysset S, von Cramon DY, et al. (2006) The Effects of Insulin on the Central Nervous System-Focus on Appetite Regulation. *Horm Metab Res* 38: 442-446.
- Rolls ET (2007) Understanding the Mechanisms of Food Intake and Obesity. *Obes Rev* 8: 67-72.
- Ribeiro G, Santos O (2013) Recompensa alimentar: mecanismos envolvidos e implicações para a obesidade. *Rev Port Endocrinol Diabetes Metab* 8: 82-88.
- D'alessio D (2008) Intestinal Hormones and Regulation of Satiety: The Case for CCK, GLP-1, PYY, and Apo A-IV. *J Parenter Enteral Nutr* 32: 567-568.
- Boey D, Heilbronn L, Sainsbury A, Laybutt R, Kriketos A, et al. (2006) Low Serum PYY is Linked to Insulin Resistance in First Degree Relatives of Subjects with Type 2 Diabetes. *Neuropeptides* 40: 317-324.
- Ukkola OH, Puurunen VP, Piira OP, Niva JT, Lepojärvi ES, et al. (2011)

- High Serum Fasting Peptide YY (3-36) is Associated with Obesity-Associated Insulin Resistance and Type 2 Diabetes. *Regul Pept* 170: 38-42.
34. Steinert RE, Feinle-Bisset C, Geary N, Beglinger C (2013) Digestive Physiology of the Pig Symposium: Secretion of Gastrointestinal Hormones and Eating Control. *J Anim Sci* 91: 1963-1973.
 35. Brandão PP, Garcia-Souza EP, Neves FA, dos Santos Pereira MJ, Sichieri R, et al. (2011) Appetite-Related Hormone Levels in Obese Women with and Without Binge Eating Behavior. *Rev Nutr* 24: 667-677.
 36. Sayegh AI (2008) PYY Decreases Food Intake and Activates Peripheral and Central Neurons in Rats. *Appetite* 51: 397.
 37. Valassi E, Scacchi M, Cavagnini F (2008) Neuroendocrine Control of Food Intake. *Nutr Metab Cardiovasc Dis* 18: 158-168.
 38. Batterham RL, Cowley MA, Small CJ, Herzog H, Cohen MA, et al. (2002) Gut Hormone PYY(3-36) Physiologically Inhibits Food Intake. *Nature* 418: 650-654.
 39. Ekblad E, Sundler F (2002) Distribution of Pancreatic Polypeptide and Peptide YY. *Peptides* 23: 251-261.
 40. Simonian HP, Kresge KM, Boden GH, Parkman HP (2005) Differential Effects of Sham Feeding and Meal Ingestion on Ghrelin and Pancreatic Polypeptide Levels: Evidence for Vagal Efferent Stimulation Mediating Ghrelin Release. *Neurogastroenterol Motil* 17: 348-354.
 41. Kojima S, Ueno N, Asakawa A, Sagiyama K, Naruo T, et al. (2007) A Role for Pancreatic Polypeptide in Feeding and Body Weight Regulation. *Peptides* 28: 459-463.
 42. Guyenet SJ, Schwartz MW (2012) Clinical Review: Regulation of Food Intake, Energy Balance, and Body Fat Mass: Implications for the Pathogenesis and Treatment of Obesity. *J Clin Endocrinol Metab* 97: 745-755.
 43. Holzer P, Reichmann F, Farzi A (2012) Neuropeptide Y, Peptide YY and Pancreatic Polypeptide in the Gut-Brain Axis. *Neuropeptides* 46: 261-274.
 44. Edelsbrunner ME, Herzog H, Holzer P (2009) Evidence from Knockout Mice that Peptide YY and Neuropeptide Y Enforce Murine Locomotion, Exploration and Ingestive behaviour in a Circadian Cycle-and Gender-Dependent Manner. *Behav Brain Res* 203: 97-107.
 45. Berntson GG, Zipf WB, O'Dorisio TM, Hoffman JA, Chance RE (1993) Pancreatic Polypeptide Infusions Reduce Food Intake in Prader-Willi Syndrome. *Peptides* 14: 497-503.
 46. Naessén S, Carlström K, Holst JJ, Hellström PM, Hirschberg AL (2011) Women with Bulimia Nervosa Exhibit Attenuated Secretion of Glucagon-Like Peptide 1, Pancreatic Polypeptide, and Insulin in Response to a Meal. *Am J Clin Nutr* 94: 967-972.
 47. Wittamer V, Grégoire F, Robberecht P, Vassart G, Communi D, et al. (2004) The C-Terminal Nonapeptide of Mature Chemerin Activates the Chemerin Receptor with Low Nanomolar Potency. *J Biol Chem* 279: 9956-9962.
 48. Bozaoglu K, Bolton K, McMillan J, Zimmet P, Jowett J, et al. (2007) Chemerin is a Novel Adipokine Associated with Obesity and Metabolic Syndrome. *Endocrinology* 148: 4687-4694.
 49. Bozaoglu K, Segal D, Shields KA, Cummings N, Curran JE, et al. (2009) Chemerin is Associated with Metabolic Syndrome Phenotypes in a Mexican-American Population. *J Clin Endocrinol Metab* 94: 3085-3088.
 50. Schmid A, Bala M, Leszczak S, Ober I, Buechler C, et al. (2016) Pro-Inflammatory Chemokines CCL2, Chemerin, IP-10 and RANTES in Human Serum During an Oral Lipid Tolerance Test. *Cytokine* 80: 56-63.
 51. Rittner HL, Stein C (2005) Involvement of Cytokines, Chemokines and Adhesion Molecules in Opioid Analgesia. *Eur J Pain* 9: 109-112.
 52. Grandison L, Guidotti A (1977) Stimulation of Food Intake by Muscimol and Beta Endorphin. *Neuropharmacol* 16: 533-536.
 53. Sanger DJ, McCarthy PS (1981) Increased Food and Water Intake Produced in Rats by Opiate Receptor Agonists. *Psychopharmacology* 74: 217-220.
 54. Morley JE, Levine AS, Murray SS, Kneip J (1982) Peptidergic Regulation of Norepinephrine Induced Feeding. *Pharmacol Biochem Behav* 16: 225-228.
 55. Papadouka V, Carr KD (1994) The Role of Multiple Opioid Receptors in The Maintenance of Stimulation-Induced Feeding. *Brain Res* 639: 42-48.
 56. Bertolini A, Tacchi R, Vergoni AV (2009) Brain effects of melanocortins. *Pharmacol Res* 59: 13-47.
 57. Panaro BL, Tough IR, Engelstoft MS, Matthews RT, Digby GJ, et al. (2014) The Melanocortin-4 Receptor is Expressed in Enterendocrine L Cells and Regulates the Release of Peptide YY and Glucagon-Like Peptide 1 *in vivo*. *Cell Metab* 20: 1018-1029.
 58. Xu B, Goulding EH, Zang K, Cepoi D, Cone RD, et al. (2003) Brain-Derived Neurotrophic Factor Regulates Energy Balance Downstream of Melanocortin-4 Receptor. *Nat Neurosci* 6: 736-742.
 59. Krude H, Biebermann H, Luck W, Horn R, Brabant G, et al. (1998) Severe Early-Onset Obesity, Adrenal Insufficiency and Red Hair Pigmentation Caused by Pomc Mutations in Humans. *Nat Genet* 19: 155-157.
 60. Hoch M, Eberle AN, Wagner U, Bussmann C, Peters T, et al. (2007) Expression and Localization of Melanocortin-1 Receptor in Human Adipose Tissues of Severely Obese Patients. *Obesity* 15: 40-49.
 61. Brito MN, Brito NA, Baro DJ, Song CK, Bartness TJ (2007) Differential Activation of the Sympathetic Innervation of Adipose Tissues by Melanocortin Receptor Stimulation. *Endocrinology* 148: 5339-5347.
 62. Rossi J, Balthasar N, Olson D, Scott M, Berglund E, et al. (2011) Melanocortin-4 Receptors Expressed by Cholinergic Neurons Regulate Energy Balance and Glucose Homeostasis. *Cell Metab* 13: 195-204.
 63. Williams DL, Bowers RR, Bartness TJ, Kaplan JM, Grill HJ (2003) Brainstem Melanocortin 3/4 Receptor Stimulation Increases Uncoupling Protein Gene Expression in Brown Fat. *Endocrinology* 144: 4692-4697.
 64. Obici S, Feng Z, Tan J, Liu L, Karkanias G, et al. (2001) Central Melanocortin Receptors Regulate Insulin Action. *J Clin Invest* 108: 1079-1085.
 65. Bazzano LA, Serdula MK, Liu S (2003) Dietary Intake of Fruits and Vegetables and Risk of Cardiovascular Disease. *Curr Atheroscler Rep* 5: 492-499.
 66. Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB, et al. (2006) Fruit and Vegetable Intakes, C-Reactive Protein, and The Metabolic Syndrome. *Am J Clin Nutr* 84: 1489-1497.
 67. Heidemann C, Schulze MB, Franco OH, van Dam RM, Mantzoros CS, et al. (2008) Dietary Patterns and Risk of Mortality from Cardiovascular Disease, Cancer, and All Causes in a Prospective Cohort of Women. *Circulation* 118: 230-237.

68. Barbosa KBF, Costa NMB, Alfenas RCG, De Paula SO, Minim VPR, et al. (2010) Oxidative Stress: Concept, Implications and Modulating Factors. *Rev Nutr* 23: 629-643.
69. Chuang C, McIntosh MK (2011) Potential Mechanisms by which Polyphenol-Rich Grapes Prevent Obesity- Mediated Inflammation and Metabolic Diseases. *Annu Rev Nutr* 31: 155-176.
70. Scalbert A, Williamson G (2000) Dietary Intake and Bioavailability of Polyphenols. *J Nutr* 130: 2073S-2085S.
71. Suganuma M, Okabe S, Oniyama M, Tada Y, Ito H, et al. (1998) Wide Distribution of [3H]-(-)-Epigallocatechin gallate, A Cancer Preventive Tea Polyphenol, in Mouse Tissue. *Carcinogenesis* 19: 1771-1776.
72. Vitrac X, Desmoulière A, Brouillaud B, Krisa S, Deffieux G, et al. (2003) Distribution of [14C]-Trans-Resveratrol, A Cancer Chemopreventive Polyphenol, in Mouse Tissues After Oral Administration. *Life Sci* 72: 2219-2233.
73. Janle EM, Lila MA, Grannan M, Wood L, Higgins A, et al. (2010) Pharmacokinetics and Tissue Distribution of 14C-Labeled Grape Polyphenols in the Periphery and the Central Nervous System Following Oral Administration. *J Med Food* 13: 926-933.
74. Ferruzzi MG, Lobo JK, Janle EM, Cooper B, Simon JE, et al. (2009) Bioavailability of Gallic Acid and Catechins From Grape Seed Polyphenol Extract is Improved by Repeated Dosing in Rats: Implications for Treatment in Alzheimer's Disease. *J Alzheimers Dis* 18: 113-124.
75. Dulloo AG, Duret C, Rohrer D, Girardier L, Mensi N, et al. (1999) Efficacy of A Green Tea Extract Rich in Catechin Polyphenols and Caffeine in Increasing 24-H Energy Expenditure and Fat Oxidation in Humans. *Am J Clin Nutr* 70: 1040-1045.
76. Park EY, Kim EH, Kim MH, Seo YW, Lee JI, et al. (2012) Polyphenol-Rich Fraction of Brown Alga *Ecklonia cava* Collected From Gijang, Korea, Reduces Obesity and Glucose Levels in High-Fat Diet-Induced Obese Mice. *Evid Based Complement Alternat Med* 2012: 1-11.
77. Lu C, Zhu W, Shen CL, Gao W (2012) Green Tea Polyphenols Reduce Body Weight in Rats by Modulating Obesity-Related Genes. *PLoS One* 7: e38332.
78. Tom FA, Andres-Lacueva C (2012) Polyphenols and Health: Current State and Progress. *J Agric Food Chem* 60: 8773-8775.
79. Gomes SF, Silva FC, Volp ACP (2015) Prospects about the Use of Açai (*Euterpe oleracea Mart.*) on the Modulation of the Inflammation and Energetic Metabolism in Overweight and Obese Individuals. *J Obesity* 1: 1-6.
80. Kiran SP (2013) Effects of Dietary Polyphenols on Neuroregulatory Factors and Pathways that Mediate Food Intake and Energy Regulation in Obesity. *Mol Nutr Food Res* 57: 34-47.
81. Weickert MO, Reimann M, Otto B, Hall WL, Vafeiadou K, et al. (2006) Soy Isoflavones Increase Preprandial Peptide YY (PYY), But Have No Effect on Ghrelin and Body Weight in Healthy Postmenopausal Women. *J Negat Results Biomed* 5: 1-11.
82. Goldstone AP (2006) The Hypothalamus, Hormones, and Hunger: Alterations in Human Obesity and Illness. *Prog Brain Res* 153: 57-73.
83. Alkemade A, Yi CX, Pei L, Harakalova M, Swaab DF, et al. (2012) AgRP and NPY Expression in the Human Hypothalamic Infundibular Nucleus Correlate with Body Mass Index, Whereas Changes in α MSH are Related to Type 2 Diabetes. *J Clin Endocrinol Metab* 97: E925-E933.
84. Lee YM, Yoon Y, Yoon H, Park HM, Song S, et al. (2017) Dietary Anthocyanins Against Obesity and Inflammation. *Nutrients* 9: E1089.
85. Yamashita AS, Belchior T, Lira FS, Bishop NC, Wesner B, et al. (2018) Regulation of Metabolic Disease-Associated Inflammation by Nutrient Sensors. *Mediators Inflamm* 2018: 1-18.