

Pediatrics and Neonatal Nursing: Open Access

Short Communication

Volume: 2.3

Open Access

Revisiting Sound in the NICU: Implications for the Developmental Timing, Amount and Type of Sound

Amanda C DeArmond*, Jennifer R Yello, Khlood S Bubshait and Charlene A Krueger

College of Nursing, University of Florida, Gainesville, Florida, USA

*Corresponding author: Amanda C DeArmond, College of Nursing, University of Florida, Gainesville, Florida, USA, E-mail: adearmond@ufl.edu

The purpose of this short communication is to invite nurses to revisit the implications for exposure to sound in the preterm infant. Since 1997, the US Environmental Protection Agency has warned that a sound level exceeding 45 dB is best avoided in the Neonatal Intensive Care Unit [1]. More recently, the seventh edition of the recommended standards for ICU design states that, "the combination of continuous background sound and operational sound shall not exceed an hourly Leq of 45 dB and an hourly L10 of 50 dB, both A-weighted slow response in infant rooms and adult sleep areas. Transient sounds or Lmax shall not exceed 65 dB, A-weighted, slow response in these rooms/areas" [2]. The Leq measures a steady (dBA) sound level across a 30 second time period, while the L₁₀ is a measure of the decibel level exceeded for 10% of the hour. The Lmax is the highest decibel level measured for at least 1/20th of a second during the hour [3,4].

Unfortunately, it has been found repetitively that the sound levels in Neonatal Intensive Care Units (NICU) continue to remain above recommended decibel levels. In 2005, Krueger et al. [5] found that the overall average hourly Leq mean in the NICU was 60.44 dB with a range of 55-68 dB, the L₁₀ had a mean of 59.26 dB and the Lmax average was 78.39 with a range of 69-93 dB. More recently, Chow and Stellhass [6] found that the average NICU sound level during the day had a mean of 56.1 ± 2.6 dB with a peak noise level of 85.8 ± 0.7 dB. Night shift had a similar mean of 54.7 ± 4.2 dB; p=0.06 with a peak noise level of 85.0 ± 1.3 dB; p<0.001.

As technology advances, more and more infants are exposed to altered auditory stimulation at still younger ages and during a time period when the auditory system is reaching structural and functional maturity (from 25 weeks' gestation to 5 to 6 months of age). For example, premature infants are placed in incubators, which are a source of increased decibel levels in the NICU, especially when the incubator doors are closed and the fans turned on [7]. Further, infants requiring respiratory support experience still higher sound levels associated with the ventilator equipment [8]. For the fetus, the time period prior to birth is when different processes become tuned to receive qualitatively different signals of specific frequencies and intensities from outside auditory stimulation [9]. In utero, the fetus hears primarily through bone conduction and through fluid while preterm infants (born <37 weeks of gestation) must hear primarily through air conduction.

The effect of these consistently elevated sound levels in the NICUs on preterm infants is staggering. Elevated sound levels have been shown to cause immediate physiological changes, such as an increase in heart rate, blood pressure and respiratory rates, and a decrease in oxygen saturation [10]. Long term, elevated sound levels can potentially result in hearing loss and delays in language development [11].

Received date: 23 May 2016; Accepted date: 07 Jul 2016; Published date: 13 Jul 2016.

Citation: DeArmond AC, Yello JR, Bubshait KS, Krueger CA (2016) Revisiting Sound in the NICU: Implications for the Developmental Timing, Amount and Type of Sound. *Pediatr Neonatal Nurs Open Access* 2(3): doi <http://dx.doi.org/10.16966/2470-0983.116>

Copyright: © 2016 DeArmond AC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Not all sound exposure within the NICU however should be considered noise or non-beneficial. A mother's voice, either spoken or sung, has been shown to be beneficial for late preterm hospitalized infants [12-14]. Moreover, live maternal voice resulted in significant decreases in critical events and increases in oxygen saturation [15].

In conclusion, nurses are well positioned to inform researchers and practitioners involved in the care of the preterm infant. It is therefore imperative that nurses revisit sound in the NICU by not only considering decibel levels but, the developmental timing, amount and type of auditory stimulation [16,17].

References

1. Etzel RA, Balk SJ, Bearer CF, Miller MD, Shea KM, et al. (1997) Noise: a hazard for the fetus and newborn. American Academy of Pediatrics. Committee on Environmental Health. *Pediatrics* 100: 724-727.
2. White RD (2007) Recommended standards for the newborn ICU. *J Perinatol* 27: S4-S19.
3. Philbin MK, Robertson H, Hall JW (1999) Recommended permissible noise criteria for occupied, newly constructed, or renovated hospital nurseries. The Sound Study Group of the National Resource Center. *J Perinatol* 19: 559-563.
4. Philbin MK, Gray L (2002) Changing levels of quiet in an intensive care nursery. *J Perinatol* 22: 455-460.
5. Krueger C, Wall S, Parker L, Nealis R (2005) Elevated sound levels within a busy NICU. *Neonatal Netw* 24: 33-37.
6. Chow VY, Shellhaas RA (2016) Acoustic environment profile of the neonatal intensive care unit: High ambient noise and limited language exposure. *J Neonatal Nurs* 22: 159-162.
7. Marik PE, Fuller C, Levitov A, Moll E (2012) Neonatal incubators: A toxic sound environment for the preterm infant?. *Pediatr Crit Care Med* 13: 685-689.
8. Shoemark H, Harcourt E, Arnup SJ, Hunt RW (2016) Characterising the ambient sound environment for infants in intensive care wards. *J Paediatr Child Health* 52: 436-440.
9. Graven SN, Browne JV (2008) Auditory development in the fetus and infant. *Newborn Infant Nurs Rev* 8: 187-193.
10. Wachman EM, Lahav A (2010) The effects of noise on preterm infants in the NICU. *Arch Dis Child Fetal Neonatal Ed* 96: F305-F309.
11. Lahav A, Skoe E (2013) An acoustic gap between the NICU and womb: a potential risk for compromised neuroplasticity of the auditory system in preterm infants. *Front Neurosci* 8: 381.

12. Dorn F, Wirth L, Gorbey S, Wege M, Zemlin M, et al. (2014) Influence of acoustic stimulation on the circadian and ultradian rhythm of premature infants. *Chronobiol Int* 31: 1062-1074.
13. Piccolini O, Porro M, Meazza A, Gianni ML, Rivoli C, et al. (2014) Early exposure to maternal voice: effects on preterm infants development. *Early Hum Dev* 90: 287-292.
14. Chorna OD, Slaughter JC, Wang L, Stark AR, Maitre NL (2014) A pacifier-activated music player with mother's voice improves oral feeding in preterm infants. *Pediatrics* 133: 462-468.
15. Filippa M, Devouche E, Arioni C, Imberty M, Gratier M (2013) Live maternal speech and singing have beneficial effects on hospitalized preterm infants. *Acta Paediatr* 102: 1017-1020.
16. Lickliter R (2000) The role of sensory stimulation in perinatal development: insights from comparative research for care of the high-risk infant. *J Dev Behav Pediatr* 21: 437-447.
17. Cardin A, Rens L, Stewart S, Danner-Bowman K, McCarley R, et al. (2015) Neuroprotective Core Measures 1–7: A Developmental Care Journey: Transformations in NICU Design and Caregiving Attitudes. *Newborn Infant Nurs Rev* 15: 132-141.