

Comparison of the Effects of Diclofenac Sodium or Celecoxib on the Healing Tooth Socket. A Study in Rats

Samira Hajisadeghi¹, Farimah Sardari², Mostafa Sadeghi³, Sayyed Mohammad Razavi⁴ and Michael W Roberts^{5*}

¹Assistant Professor, Dental and Oral Research Center, Department of Oral and Maxillofacial Medicine, School of Dentistry, Qom University of Medical Sciences, Qom, Iran

²Assistant Professor, Department of Oral and Maxillofacial Medicine, School of Dentistry, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

³Professor, Department of Operative Dentistry, School of Dentistry, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

⁴Professor, Department of Oral and Maxillofacial Pathology and Torabinejad Dental Research center, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

⁵Professor, Department of Pediatric Dentistry, University of North Carolina School of Dentistry, Chapel Hill, North Carolina, USA

*Corresponding author: Michael W Roberts, Professor, Department of Pediatric Dentistry, University of North Carolina School of Dentistry, Chapel Hill, North Carolina, USA, Tel: (919)537-3781; E-mail: Mike_Roberts@unc.edu

Received: 26 Jun, 2018 | Accepted: 04 Jul, 2018 | Published: 10 Jul, 2018

Citation: Hajisadeghi S, Sardari F, Sadeghi M, Razavi SM, Roberts MW (2018) Comparison of the Effects of Diclofenac Sodium or Celecoxib on the Healing Tooth Socket. A Study in Rats. *J Surg Open Access* 4(2): dx.doi.org/10.16966/2470-0991.170

Copyright: © 2018 Hajisadeghi S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: There is extensive evidence in the orthopedic literature that prolonged use of Nonsteroidal Anti-Inflammatory Drugs (NSAIDs) can hinder long bone fracture healing and new bone formation around implants. The aim of this study was to investigate if short time administrations of NSAIDs such as diclofenac or celecoxib interfere with the course of healing of alveolar bony socket following tooth extraction.

Materials and Methods: Forty-five rats (15 per group) were used. After extraction of the right maxillary first molar, 15 rats received 5 mg/kg/day of diclofenac, 15 rats received 15 mg/kg/day of celecoxib and 15 rats received normal saline. The animals were sacrificed at 7, 14 and 21 days following tooth extraction. The number of osteoclasts, osteoblast and new bone formation were determined by using histological analyses. Data were analyzed by one-way ANOVA followed by Tukey's post hoc test ($\alpha=0.05$).

Results: On the 7th day, the osteoblast number in the control group was higher than the diclofenac and celecoxib groups. Tartrate-Resistant Acid Phosphatase (TRAP) immunolabeling of the control group was more than the diclofenac group on the 7th day and more than the celecoxib group on the 14th day. On the 21st day, no significant differences were noted among the three studied groups. On the 14th and 21st day, new bone formation in the NSAID treated rats was not significantly different from that in the control rats.

Conclusion: Short-term treatment with diclofenac or celecoxib, although they have the capacity to inhibit the enzyme cyclooxygenase, did not cause a significant delay in alveolar bone healing during the experimental period in rats.

Keywords: Anti-inflammatory agents; Celecoxib; Diclofenac; NSAIDs; Alveolar bone; Healing

Introduction

The conversion of arachidonic acid to prostaglandins was catalyzed by the cyclooxygenase enzymes COX-1 and COX-2, which modulate osteoblastic and osteoclastic functions and exert an anabolic effect on bone in the healing process [1]. The nonsteroidal anti-inflammatory drugs (NSAIDs) inhibit COX-1 and COX-2 activity and, consequently, prostaglandins synthesis [2]. NSAIDs are frequently used in the control of fever and pain, edema, in acute post-surgical or post-traumatic pain, and for relief of chronic pain associated with musculoskeletal disorders. NSAIDs are commonly used drugs for management

of pain after tooth extraction [3]. The NSAIDs may delay the repair of damaged tissues, such as the fractured bone, cartilage, skin and new bone formation after tooth extraction [4,5].

It has been shown in rodent studies that NSAIDs may disturb not only bone healing but also normal bone growth [6,7]. Despite the unequivocal evidence that NSAIDs may delay recovery of damaged tissues, the precise mechanism of this effect is not fully understood. Currently, numerous NSAIDs are available, from the conventional nonselective NSAIDs which inhibit non-selectively both COX-1/COX-2 enzymes, toward a new class of highly selective COX-2

inhibitors, the coxibs. Diclofenac (inhibits non-selectively both COX-1 and COX-2 enzymes) and celecoxib (selectively inhibits COX-2) are NSAIDs presenting significant analgesic and anti-inflammatory activities [3,8].

In orthopedics, prolonged use of both nonselective and selective NSAIDs can hinder repair long bone formation [8]. Even though, the conventional NSAIDs are commonly used in dentistry for pain and swelling management, very few assessments have been conducted to assess their deleterious effects on healing of the alveolar bony socket [9].

There are only limited data on the effects of NSAIDs on wound healing after oral surgery. The present study was designed to investigate whether diclofenac and celecoxib have a detrimental effect on alveolar bone healing, taking the filling of extraction socket with new formed bone as an experimental model of alveolar bone formation.

Materials and Methods

Forty five male Wistar rats at 8-10 weeks of age, weighing 200 ± 25 g were used in this study. The animals were housed under similar conditions (22°C room temperature, 40% humidity and 12 hours light/12 hours dark cycle, with free access to water and rat chow) during the study period. This study was assessed and approved by Animal Research Ethics Committee of the Isfahan University of Medical Sciences, Iran and the principles of laboratory animal care and national laws on animal use were followed in the present study [10].

General anesthesia was induced by intramuscular injection of ketamine 10% (Alfasan International, Woerden, Holland, 80 mg/kg) and Xylazine (Neurotranq, Alfasan, Woerden, Holland, 8 mg/kg) in all rats and their right upper molars were luxated with the aid of a tapered instrument and extracted with a modified hemostat (two cavities were made in each beak). An atraumatic surgical technique was used to extract the teeth. The animals were observed until fully recovered; no antibiotic or other medication was used.

The animals were randomly assigned into three groups. The celecoxib group received a daily dose (15 mg/kg body weight of celecoxib (Daroupakhsh Co, Tehran, Iran) by gavage administration; the diclofenac group received a daily dose (5 mg/kg body weight of diclofenac sodium (Daroupakhsh Co) diluted with sterile distilled water and was injected subcutaneously. The control group received only normal saline, daily. Medications in all groups were administered for a period of seven days, starting on the day of tooth extraction. Doses of all drugs were chosen based on prior studies and pharmacokinetic data to mimic similar doses used to treat humans, taking into account the differences in metabolism between species [3,11,12].

Five animals from each group were sacrificed on the 7th, 14th and 21st day post extraction by an over dose of ether inhalation. The mandibles was removed and the maxillas were fixed in 4% paraformaldehyde solution (SRPC, Tehran, Iran), demineralized with 10% Ethylenediaminetetraacetic acid (EDTA) (Merck, Darmstadt, Germany) and embedded in paraffin (Merck). The pieces were sectioned perpendicular to the long axis of the alveolar process with a microtome (Accu-Cut SRM, SAKURA, USA) in order to obtain slices of five μm thicknesses, which were mounted in previously poly-L-lysine slides.

A slide using Hematoxylin-Eosin (H&E) stain was prepared from each specimen. Sections representative of the number of osteoblast and amount of new bone formation in each tooth were captured by a digital camera (Axio Cam MRC5; Carl Zeiss do Brasil Ltda, Rio de

Janeiro, RJ, Brazil) coupled to a light microscope (Olympus CX21FS, Olympus Corporation, Tokyo, Japan) with 1:100 magnification. For the immunohistochemistry reactions, blocking with 0.03% hydrogen peroxide followed by primary antibodies anti Tartrate-Resistant Acid Phosphatase (TRAP) (Goat anti trap polyclonal-Santa Cruz, CA, USA) and the biotinylated donkey anti-goat antibodies (Biotin-SP-AffiniPure donkey anti-goat IgG-Jackson Immunoresearch Laboratories, West Grove, PA, USA) was the secondary antibody; the immunohistochemistry reaction signal was amplified with the Avidin-Biotin system (Kit ABC Vectastain Elite ABC, Vector Laboratories, Burlingame, CA, USA) and the reaction was revealed using diaminobenzidine (DAB-Sigma, Saint Louis, MO, USA) as the chromogen.

Additional sections were used for immunohistochemical staining to determine the expression of TRAP protein in the alveolar tissues during the healing process from tooth extraction, and then were counterstained with Harris's hematoxylin. Immunostaining was used to evaluate alveolar bone under conventional optical microscope. A negative control was prepared for each specimen using the same method except for the primary antibody. Sections representative of TRAP protein in each tooth were captured by a digital camera (Axio Cam MRC5; Carl Zeiss do Brasil Ltda., Rio de Janeiro, RJ, Brazil) coupled to a light microscope (Olympus CX21FS, Olympus Corporation, Tokyo, Japan) with 1:400 magnification. Slides were examined by an expert blinded pathologist.

The results are presented as means \pm SD, and statistical differences among studied groups were assessed by one-way ANOVA followed by Tukey's post hoc test. All analyses were done using SPSS-20 (SPSS Inc., Chicago, IL, USA) and P value less than 0.05 were considered significant.

Results

Throughout the study period, all animals tolerated the experiment. Behavior and body weight among the groups were not significantly different ($P > 0.05$).

The results of the comparison of osteoblast number, TRAP (osteoclast number) and new bone formation in the dental socket among the groups are shown in figures 1 and 2. As shown, the number of osteoblast on the 7th day in the control group was significantly greater than the diclofenac and celecoxib groups ($P < 0.05$); but at 14th and 21st day, differences among the groups were not statistically different ($P > 0.05$) (Figure 1). The number of TRAP-positive osteoclasts among the groups at 7 and 14 days were statistically significant ($P < 0.05$). The control number of TRAP was significantly higher than the diclofenac and the celecoxib groups at 14 days ($P < 0.05$). Although, the number of osteoclasts observed in the control group was greater compared with diclofenac and celecoxib groups on the 21st day, the differences were not statistically significant ($P > 0.05$) (Figure 3). New bone formation at 7 days among the studied groups was significantly different. The new bone growth in the diclofenac and celecoxib groups was significantly less than the control group at 7 days ($P < 0.05$), but the difference between the diclofenac and celecoxib groups was not significant ($P > 0.05$). On the 14th and 21st day, new bone formation among studied groups was similar and there were no statistical significant among groups ($P > 0.05$) (Figures 3 and 4).

Discussion

A suitable model for the study of bone formation is provided for alveolar healing and can be considered a sensitive indicator of bone damage and repair [13,14]. In different animal species, the process of




Figure 1: The number of osteoblasts in the three groups.




Figure 2: New bone formation in the three groups.




Figure 3: The number of TRAP-positive osteoclasts in the three groups.

intra-alveolar healing is well established [15,16]. In brief, after a tooth is extracted the socket is filled with a blood clot which is progressively invaded by capillary sprouting and fibroblasts originating from periodontal ligament remnants. In the sequence, the amount of inflammatory cells and blood vessels decreases, the osteoblasts become evident, and the coagulum is gradually absorbed and replaced by immature connective tissue. An immature bone matrix initially




Figure 4: New bone formation at 7th, 14th and 21st post-extraction days in the control, diclofenac and celecoxib groups.

synthesize by osteoblasts that is further mineralized by calcium deposition as hydroxyapatite crystals. The alveolar bone neof ormation takes place toward the center from the lateral and apical walls, and the healing process culminates with filling of the dental socket by trabecular bone [17,18].

In this study, the histological feature of the alveolar socket healing was observed in a control and animals treated with two different types of NSAIDs (diclofenac and celecoxib) and followed for 21 days. It has been shown in previous rat studies that the maximum mineral bone density and the major proportion of bone formation takes place by the end of the second week after tooth extraction [19,20]. In this study the hypothesis that NSAIDs could hinder the alveolar bone healing was tested. Our results showed that in the first week osteoblast, TRAP-positive osteoclasts and new bone formation of diclofenac and celecoxib groups were lower than the control group. The differences between the control group and the study groups in osteoblast, osteoclast and bone formation was significant in the first week, there was no significant difference between the three groups in these criteria by the end of the third week. It would appear that the short-time administration of a low dose of NSAID for pain relief after tooth extraction may cause initial delay in healing, but no important negative effect occurred in long term bone healing. *In vivo* studies found similar impairments of fracture healing by NSAIDs, primarily in the early phase of healing [21,22].

Osteoclasts are distinguished from macrophages by the presence of TRAP in their cytoplasm. This type-V isoenzyme of acid phosphatase presents an intense activity in osteoclasts, being considered a specific marker for osteoclasts. Histological analysis is an effective method to identifying osteoclasts and analyzing their dynamics as is staining of TRAP, a cytochemical marker for osteoclasts [23]. In our study, diclofenac sodium had lower clastic activity, as demonstrated by lower TRAP expression. This drug also led to a decrease in TRAP expression three weeks post extraction; especially in the first seven days. Histologically observations also confirmed the presence of clastic cells. Celecoxib led to lower clastic activity, especially in the second week. Other studies have shown that celecoxib and diclofenac reduced osteoclastogenesis *in vitro* [11,24,25]. They have also shown that NSAIDs inhibit the number of osteoblast in rats at a very early stage [26,27].

A nonselective inhibition of the enzymes COX-1 and COX-2 by conventional NSAIDs have demonstrated interference with long bone fracture healing and rate of spinal fusion but an absence of this deleterious effect has also been observed. The effects of selective COX-2 inhibition by NSAIDs were evaluated in several studies but the results were equivocal even though the importance of COX-2 and prostaglandin E2 for bone formation is well known [8].

The effect of COX inhibition on alveolar bone healing is minimal. It has also been reported that short-term treatment (four days) with diclofenac in rats delayed alveolar socket healing [28], while other studies showed that NSAIDs didn't hinder alveolar bone healing [27]. There are few studies that have used NSAIDs to control edema and pain after removal of impacted third molars and often refer to short-term drug administration. Some authors have reported no significant difference in the rate of bone healing following administration of aspirin, diclofenac, ibuprofen, and flurbiprofen [9,29,30].

The hypothetic disadvantageous effect of diclofenac and celecoxib on the healing of alveolar bony socket was not established in the present study. Some factors may explain the disagreements in the experimental outcomes about the effects of COX inhibitors on bone healing, such as duration of treatment and dosage, as well as age, intra and inter species differences regarding sensitivity to drugs, pharmacokinetics of drugs, compensatory local and systemic factors, and rate of bone remodeling [31,32].

Conclusion

The present results provide evidence that diclofenac and celecoxib do not interfere significantly with alveolar bone healing. Also, additional studies in a diversity of experimental models to definitely confirm, or discharge, the lack of interference of this drug with new bone formation with respect to duration of treatment and dosage is indicated.

Acknowledgment

We thank to the Department of Oral and Maxillofacial Medicine and Torabinejad Dental Research Center, School of Dentistry, Isfahan University of Medical Sciences, Isfahan for financial support of this study.

Conflict of Interest

The authors have no financial or personal relationship with other people or organizations that could inappropriately influence or bias this paper.

References

- Mauro A, Lipari L, Leone A, Tortorici S, Burruano F, et al. (2010) Expression of cyclooxygenase-1 and cyclooxygenase-2 in normal and pathological human oral mucosa. *Folia Histochem Cytobiol* 48: 555-563.
- Radi ZA, Khan NK (2005) Effects of cyclooxygenase inhibition on bone, tendon, and ligament healing. *Inflamm Res* 54: 358-366.
- Ghalayani P, Minaian M, Razavi SM, Hajisadeghi S, Naghsh N, et al. (2014) Effects of diclofenac and celecoxib on osteoclastogenesis during alveolar bone healing, *in vivo*. *Dent Res J* 11: 357-363.
- Pountos I, Georgouli T, Calori GM, Giannoudis PV (2012) Do Nonsteroidal Anti-Inflammatory Drugs Affect Bone Healing? A Critical Analysis. *Sci World J* 2012: 606404.
- Chikazu D, Fujikawa Y, Fujihara H, Suenaga H, Saijo H, et al. (2011) Cyclooxygenase-2 activity is important in craniofacial fracture repair. *Int J Oral Maxillofac Surg* 40: 322-336.
- Murnaghan M, Li G, Marsh DR (2006) Nonsteroidal anti-inflammatory drug-induced fracture nonunion: an inhibition of angiogenesis? *J Bone Joint Surg Am* 88: 140-147.
- Altman RD, Latta LL, Keer R, Renfree K, Hornicek FJ, et al. (1995) Effect of nonsteroidal antiinflammatory drugs on fracture healing: a laboratory study in rats. *J Orthop Trauma* 9: 392-400.
- Lamano-Carvalho TL (2007) Effect of conventional and COX-2 selective non-steroidal antiinflammatory drugs on bone healing. *Acta Ortop Bras* 15: 166-168.
- Fracon RN, Teófilo JM, Satin RB, Lamano T (2008) Prostaglandins and bone: potential risks and benefits related to the use of nonsteroidal anti-inflammatory drugs in clinical dentistry. *J Oral Sci* 50: 247-352.
- National Research Council, Division on Earth and Life Studies, Institute for Laboratory Animal Research, Committee for the Update of the Guide for the Care and Use of Laboratory Animals (2011) *Guide for the care and use of laboratory animals*. 8th Edition, National Academies Press, USA 4-15.
- Taketa T, Sakai A, Tanaka S, Nakai K, Menuki K, et al. (2008) Selective cyclooxygenase-2 inhibitor prevents reduction of trabecular bone mass in collagen-induced arthritic mice in association with suppression of RANKL/OPG ratio and IL-6 mRNA expression in synovial tissues but not in bone marrow cells. *J Bone Miner Metab* 26: 143-151.
- Bort R, Ponsoda X, Jover R, Gómez-Lechón MJ, Castell JV (1999) Diclofenac toxicity to hepatocytes: a role for drug metabolism in cell toxicity. *J Pharmacol Exp Ther* 288: 65-72.
- Teófilo JM, Giovanini GS, Fracon RN, Lamano T (2011) Histometric study of alveolar bone healing in rats treated with the nonsteroidal anti-inflammatory drug nimesulide. *Implant Dent* 20: e7-e13.
- Luvizuto ER, Queiroz TP, Dias SMD, Okamoto T, Dornelles RC, et al. (2010) Histomorphometric analysis and immunolocalization of RANKL and OPG during the alveolar healing process in female ovariectomized rats treated with oestrogen or raloxifene. *Arch Oral Biol* 55: 52-59.
- Vieira AE, Repeke CE, Ferreira Junior Sde B, Colavite PM, Biguetti CC, et al. (2015) Intramembranous bone healing process subsequent to tooth extraction in mice: micro-computed tomography, histomorphometric and molecular characterization. *PLoS One* 10: e0128021.
- Calixto RF, Teófilo JM, Brentegani LG, Lamano-Carvalho TL (2007) Alveolar wound healing after implantation with a pool of commercially available bovine bone morphogenetic proteins (BMPs): a histometric study in rats. *Braz Den J* 18: 29-33.
- Elsubeihi ES, Heersche JN (2004) Quantitative assessment of post-extraction healing and alveolar ridge remodelling of the mandible in female rats. *Arch Oral Biol* 49: 401-412.
- Shahabooei M, Razavi SM, Minaian M, Birang R, Behfarnia P, et al. (2015) A histomorphometric study of the effect of doxycycline and erythromycin on bone formation in dental alveolar socket of rat. *Adv Biomed Res* 4: 71.
- Mavropoulos A, Ödman A, Ammann P, Kiliaridis S (2010) Rehabilitation of masticatory function improves the alveolar bone architecture of the mandible in adult rats. *Bone* 47: 687-692.
- Mavropoulos A, Kiliaridis S, Bresin A, Ammann P (2004) Effect of different masticatory functional and mechanical demands on the structural adaptation of the mandibular alveolar bone in young growing rats. *Bone* 35: 191-197.

21. Bhandari M, Schemitsch EH (2002) Bone formation following intramedullary femoral reaming is decreased by indomethacin and antibodies to insulin-like growth factors. *J Orthop Trauma* 16: 717-722.
22. Hogevoid HE, Groggaard B, Reikeras O (1992) Effects of short-term treatment with corticosteroids and indomethacin on bone healing. A mechanical study of osteotomies in rats. *Acta Orthop Scand* 63: 607-611.
23. Cole AA, Wezeman FH (1987) Cytochemical localization of tartrate-resistant acid phosphatase, alkaline phosphatase, and nonspecific esterase in perivascular cells of cartilage canals in the developing mouse epiphysis. *Am J Anat* 180: 237-242.
24. Kawashima M, Fujikawa Y, Itonaga I, Takita C, Tsumura H (2009) The effect of selective cyclooxygenase-2 inhibitor on human osteoclast precursors to influence osteoclastogenesis *in vitro*. *Mod Rheumatol* 19: 192-198.
25. Kellinsalmi M, Parikka V, Risteli J, Hentunen T, Leskelä HV, et al. (2007) Inhibition of cyclooxygenase-2 down-regulates osteoclast and osteoblast differentiation and favours adipocyte formation *in vitro*. *Eur J Pharmacol* 572: 102-110.
26. Krischak G, Augat P, Blakytyn R, Claes L, Kinzl L, et al. (2007) The non-steroidal anti-inflammatory drug diclofenac reduces appearance of osteoblasts in bone defect healing in rats. *Arch Orthop Trauma Surg* 127: 453-458.
27. Chang JK, Wang GJ, Tsai ST, Ho ML (2005) Nonsteroidal anti-inflammatory drug effects on osteoblastic cell cycle, cytotoxicity, and cell death. *Connect Tissue Res* 46: 200-210.
28. Yugoshi LI, Sala MA, Brentegani LG, Lamano Carvalho TL (2002) Histometric study of socket healing after tooth extraction in rats treated with diclofenac. *Braz Dent J* 13: 92-96.
29. Baratieri A, Deli R (1979) The effect on bone repair of aspirin cones placed in extraction sockets in dogs: a histopathologic study. *J Oral Pathol* 8: 198-206.
30. Battisti N (1994) The evaluation of the analgesic and anti-inflammatory effects of flurbiprofen mouthwash and 100-mg tablets in oral medicine. *Minerva Stomatol* 43: 141-144.
31. Ferry ST, Dahnert LE, Afshari HM, Weinhold PS (2007) The effects of common anti-inflammatory drugs on the healing rat patellar tendon. *Am J Sports Med* 35: 1326-1333.
32. Mullis BH, Copland ST, Weinhold PS, Miclau T, Lester GE, et al. (2006) Effect of COX-2 inhibitors and non-steroidal anti-inflammatory drugs on a mouse fracture model. *Injury* 37: 827-837.